

# NHS GGC Adult, Mental Health, Paediatric and Neonatal Services

## Administration of Intravenous (IV) Medicines and Flush Policy (version 2)

<b>Lead Author:</b>	Lynne Robertson, Practice Development Nurse
<b>Lead Manager:</b>	Emma Henderson, Lead Nurse Excellence in Care
<b>Responsible Director:</b>	Prof Angela Wallace
<b>Approved by:</b>	NHSGGC Board Clinical Governance Forum Acute Services Clinical Governance Forum Mental Health Clinical Governance Forum Health and Community Partnership Clinical Governance Forum Area Drugs and Therapeutics Safer Use of Medicines Committee (ADTC)
<b>Date approved:</b>	23 <sup>rd</sup> September 2024
<b>Date for Review:</b>	2 years
<b>Replaces previous version: [if applicable]</b>	Administration of Intravenous (IV) Medicines and Flush Policy (version 1) 2021

This policy has been reviewed by the Equality and Human Rights Team with no risk of unfair impact on protected characteristics noted.

# 1. Contents page

2.	Introduction .....	3
3.	Scope .....	3
4.	Roles and responsibilities .....	4
5.	Consent .....	4
6.	Prescriptions .....	4
7.	IV medicine administration education.....	6
8.	Independent 2 person check.....	7
9.	Infection Control.....	10
10.	General information on the preparation and administration of IV medicines .....	13
11.	Labelling.....	16
12.	Priming and changing infusion systems.....	16
13.	Infusion devices and infusion charts .....	17
14.	IV flush .....	18
15.	Review of policy .....	20
16.	Appendix 1: preparation and administration of an IV bolus .....	21
17.	Appendix 2: preparation and administration of medicines via a volumetric device .....	23
18.	Appendix 3: preparation and administering medicines via a syringe device .....	27
19.	Appendix 4: Resources .....	30
	Communication and Implementation Plan .....	31
	Monitoring.....	31
	Impact Assessment.....	32

## 2. Introduction

The aim of this policy is to inform healthcare practitioners working in NHS Greater Glasgow and Clyde (NHSGGC) of their responsibilities in safe and effective intravenous (IV) medicines administration to reduce the risk of harm. This may include IV bolus, intermittent or continuous infusion administrations via a vascular access device (VAD).

For the purposes of this policy, VADs include peripheral venous catheters (PVC), midlines, all types of central venous catheter (CVCs) and implantable ports. It is the responsibility of individual practitioners to ensure that they are competent in accessing and managing VADs. This will require additional competencies to be achieved. Information on care and maintenance of VADs can be found in NHSGGC Vascular Access Devices, Care and Maintenance Guideline (2023). Further advice is available from Practice Development, the Vascular Access Service, specialist nurses and/or infection control nurses.

For the purposes of this policy all health care practitioners are accountable for maintaining competence in accordance with the code of conduct and guidance of their professional body, to ensure they have the knowledge and skills to deliver safe and effective practice (Nursing and Midwifery Council (NMC) (2018), Health and Care Professions Council (HCPC) (2016), General Medical Council (GMC) (2013).

Prescriptions must be written on the current approved prescribing documentation, or prescribed electronically, in accordance with current local procedures.

## 3. Scope

For the purpose of this policy the term practitioner will include all registered health care professionals (registered nurses/midwives/operating department practitioners (ODPs)/advanced nurse practitioners (ANPs), allied health care professionals (AHPs) and medical staff) within NHSGGC who are required to administer IV medicines (including IV flush) as part of their role. This includes bank and agency practitioners.

*However, registered health care professionals will work in clinical situations where it is not possible to adhere rigidly to an independent second check as outlined throughout this policy (such as medical staff in theatres/emergencies; Hospital@Home (H@H) or district nursing teams). In these situations, a paused single check or paused 2 person check (e.g. at H@H or district nurse base and patient home) is permissible. Individual practitioners are accountable to*

*ensure patient safety is not compromised.*

This policy should be used in conjunction with other relevant guidelines and standards referred to in Appendix 1.

Staff in specialist clinical areas caring for particularly vulnerable patient groups may have local standard operating procedures (SOP) in use which should be referred and adhered to.

## 4. Roles and responsibilities

For the administration of IV medicines, all practitioners must be appropriately trained and directly supervised until considered competent. A practitioner can be described as competent if they have had the necessary training, clinical experience, skills and knowledge to undertake a task safely. Before any practitioner can commence IV medicines administration education, they must be competent in the administration of medicines via other routes e.g. oral/subcutaneous/per rectum; and in the care of patients with a VAD in accordance with the requirements of their professional role.

## 5. Consent

Consent is required before practitioners undertake any care for a patient. This may be informal (verbal) or formal (written) for more complex procedures. If there is evidence of impaired capacity, either temporarily or permanently, an adult with incapacity form (AWI) should be completed by medical staff to allow health care practitioners to provide treatment that is required. In emergency situations, practitioners should use clinical judgement as to whether the risks of delaying a procedure outweigh the need for formal consent. Where a patient doesn't have English as a first language or requires additional communication support, this must be provided in line with [NHS GGC Interpreting, Communication, Support and Translation policy](#) (2023) The patient should be fully informed about treatment and potential side effects; and provided with patient information leaflets where necessary.

Children and young people: Those under 16 years have legal capacity to consent (or refuse) treatment on their own behalf, if they are deemed capable of understanding the nature and possible consequences of treatment. A parent or legal guardian may consent to medical treatment *if* the child lacks decision-making capacity.

## 6. Prescriptions

Medicines must be prescribed by a registered Doctor/ Dental Practitioner or a practitioner who has successfully completed a professionally recognised prescribing course to allow independent prescribing.

All prescribers will follow the guidance laid out in the NHSGGC Safe and Secure Handling of Medicines (2023).

Permitted abbreviations:

Kg = kilogram

g = gram

mg = milligram

L = litre

ml = millilitre

There are **no** permitted abbreviations for **microgram**, **nanogram** or **international unit** and these should be written in full.

Standardised NHSGGC documentation should be used where possible. [Samples of charts](#) are available on StaffNet.

Prescription records may include, but are not limited to:

- Electronic prescription record (e.g. Hospital Electronic Prescribing and Medicine Administration (HEPMA) Chemocare or CareVue)
- Main medicine prescription chart (Kardex)
- Specific medicine prescription form e.g. insulin, vancomycin or gentamicin
- Intravenous infusion fluid prescription charts
- Parenteral nutrition prescription
- Discharge prescription
- Patient Controlled Analgesia (PCA) chart
- Continuous infusion pump chart
- Patient Group Directive (PGD)

It is the responsibility of the individuals involved in the prescription and administration process to ensure that the correct prescribing and monitoring documentation has been commenced. This may include more than one of the prescription documents listed above. Adult and paediatric versions are available and there may be differences between these documents.

**Administration of medications without a written prescription:** Medicines must only be administered without a written prescription in emergency situations. A verbal instruction to

administer medicines via telephone is not acceptable, unless in exceptional circumstances and must be approved by the practitioner in charge. HEPMA can facilitate “remote” prescribing in certain clinical areas / situations. Local procedures should be followed and remote prescribing must only be used if it is in the patient’s best interests and the prescriber has all the clinical information required to make safe prescribing decisions e.g.

- the patient’s current and past health status
- clinical evidence for the medicine / treatment being prescribed
- compatibility of any medicine being prescribed with other current medicine / treatment the patient is receiving (including self-purchased / administered medication)

Appropriate documentation must be kept (e.g. in the patient’s medical notes) to support remote prescribing decisions.

## 7. IV medicine administration education

The [NHSGGC Medicine Administration Competency Assurance Record \(2023\)](#) should be utilised where all new registered nurses/midwives and ODPs starting substantive or bank contract employment within NHSGGC. Section A should be completed within 2 weeks. Section B may take longer. Section C should be completed when both Sections A & B are completed.

The administration of IV medication is carried out by practitioners who have successfully completed the NHSGGC IV medicines administration competency programme, or equivalent.

If a practitioner has completed an IV medicines administration course out with NHSGGC and has evidence of achieving competence (certificate or competency assessment documentation) they **do not** need to repeat the programme in NHSGGC. The practitioner undertakes supervised practice to demonstrate safe and effective practice using appropriate NHSGGC documentation.

If a practitioner has successfully completed an adult IV medicine administration competency programme and moves to paediatric/neonatal services, they must undertake a paediatric IV medicines administration competency programme.

The practitioner will:

Attend and complete the recognised NHSGGC (or equivalent) IV medicine administration competency programme; which includes education on:

- Professional aspects of IV medicine administration
- Infection control aspects of IV medicine administration
- Pharmaceutical aspects of IV medicine administration, including antimicrobial stewardship

- Theoretical aspect of IV medicine administration
- Calculations and a calculations assessment

Student nurses (all fields) and midwives will complete the equivalent theoretical aspects in part 3 of the pre-registration programme. The practitioner/student nurse/midwife will then undertake supervised practice and achieve competence in the preparation and administration of IV medicines in their year 3 placement area. Supervised practice will be under direct supervision of a suitably competent registered nurse/midwife and involve **peripheral route only**.

Although student nurses and midwives can be involved in the checking, preparation and administration of IV medicines and flush via peripheral route, consideration should be given to the complexity of calculations, experience of the supervising registered nurse/midwife and the available VAD. In some instances, 2 suitably prepared and competent registered staff will be appropriate. In all instances, the registered practitioner(s) has professional accountability of the process.

**Registered** practitioners who have **not** completed the NHSGGC IV medicines competency programme or equivalent:

- **Cannot** administer IV medicines or flush VADs – unless as part of supervised practice for completion of NHSGGC cannulation or IV medicine administration competency programmes
- Can be the second checker for IV medicines preparation or administration
- Can be second checker for IV controlled medicine administration
- Can be the first checker for IV fluids (e.g. 0.9% sodium chloride, 5% glucose, hartmanns) but not any fluids with additives e.g. potassium chloride

## 8. Independent 2 person check

In NHSGGC all IV medications are required to be checked by 2 competent practitioners prior to administration with ideally, *both* practitioners competent in IV medicine administration. However, it is acknowledged that in practice this may sometimes not be possible. Providing that one of the practitioners is IV trained, the second checker **can be** a competent practitioner who has not completed the NHSGGC IV medicine administration competency programme, or equivalent (e.g. part 3 student midwives and nurses).

Both practitioners checking the medication should be aware of its use, side effects, recommended route and method of administration prior to the administration of the medicine.

**Both practitioners should INDEPENDENTLY perform any calculations, check all ampoules or vials against prescription and expiry dates, preparation, patient identification process and commencement of administration of all medications (e.g. attachment of the medication to the vascular access device).**

The independent 2 person check should be undertaken at the same time. If this is not practically possible, both practitioners are responsible and accountable to ensure good communication with each other, and raise discrepancies and concerns.




As part of the independent 2 person check, both practitioners **must independently** confirm that:

- The medicine prescription/infusion prescription chart is fully completed
- The patient is not allergic to the medicine prescribed
- The correct patient has been identified and is wearing a name band
- The prescribed medicine/dose and infusion fluid is appropriate for the patient
- The medicine(s), any diluent/reconstituting fluids and flushing agents are within their expiry dates and compatible with one another and with any IV fluids being administered at the same time. If compatibility is unknown seek pharmacist and/or medical staff advice prior to preparation/administration
- The scheduled dose has not already been administered; supplementary charts/documentation/'once only' section of the prescription chart (electronic or paper) must be checked
- Independent calculations **must be** performed by each practitioner to confirm that the prescribed dose is correct and that the end volume/concentration/rate to be administered are correct. Discuss any discrepancies prior to medicine administration with the prescriber and seek advice from pharmacy if required
- The solution to be administered is free of visible particles and is of the correct colour
- Independent pump rate calculation **must** be performed by each practitioner to confirm rate is correct. The set rate on the device is checked and verified by both practitioners
- Commencement of administration by witness of connection of syringe or administration set to the vascular access device
- Changes to device rates should involve **an additional** independent 2 person check

If either practitioner is concerned or unsure about any aspect of the prescription, knowledge of medication or skills they should not administer the medicine, inform the patient's medical practitioner and seek support.

Medicines should be administered **immediately** following preparation, where practically possible. **'Chance to Check'** is a conscious safety pause immediately prior to administration, which highlights the importance of the correct administration of medications. It raises awareness of what administration practices are acceptable and uses five key statements as part of a checking process and should be undertaken as part of the independent 2 person check by both

practitioners. Chance to Check also reinforces the need for staff administering medications to take the time to ensure the task is carried out efficiently, without interruption or delay.



## 9. Infection Control

**Standard Infection Control Precautions (SICPs):** Current local and [national guidance](#) advise that SICPs should be embedded into all aspects of care delivery including care of patients receiving IV medicines. The principles of SICPs should be used **by all staff in all care settings at all times for all patients** whether infection is known to be present or not. The application of SICPs ensures the safety of patients, staff and visitors and is determined by the degree of risk

encountered including the task/level of interaction and/or the level of exposure to blood or other body fluids.

There are 10 elements which make up SICPs, 6 of which must be applied for all procedures within this policy:

**Hand hygiene:** Practitioners must undertake [hand hygiene at key moments](#) to reduce the risk of contamination and infection.

**Choice of Personal Protective Equipment (PPE):** Before undertaking any procedure, practitioners must assess any likely exposure to blood and/or body fluids and ensure that PPE is worn to provide protection for the practitioner. A minimum of disposable purple apron and non-sterile nitrile gloves must be worn. Gloves should be changed if they become visibly contaminated and/or a perforation/puncture is suspected. In addition, staff should consider the need for a surgical mask/eye protection if splashing of blood or body fluids is anticipated. [Control Of Substances Hazardous to Health \(COSHH\) policy](#) should also be adhered to in all aspects of the preparation and administration process. All PPE must be removed on completion of task and immediately placed in a clinical healthcare waste bin after which hand hygiene **must** be undertaken.

Additional precautions may be necessary for different types of VADs e.g. sterile gloves may be required if there is risk of contamination of sterile key parts/sites as part of an aseptic non touch technique (ANTT®). It is the practitioner's responsibility to undertake a risk assessment and choose appropriate equipment for personal and patient protection. **Aseptic Non Touch Technique (ANTT®):** A non touch technique should be adopted when administering IV medications to maintain asepsis. This involves not touching the key parts/sites such as the tip of a syringe or a needle free access device/access port once it has been decontaminated. The needle free access device or access port must be cleaned with 2% chlorhexidine in 70% isopropyl alcohol for 30 seconds and allowed to dry before administering IV medicines/flush. The risk of contamination of these key parts/sites must be assessed to choose the appropriate level of precaution. For example, non sterile gloves are acceptable in the majority of circumstances when accessing VADs using a non touch technique, however, sterile gloves may be chosen in more complex procedures to reduce risk of contamination. The clinician should risk assess each procedure, bearing in mind the condition and location of the patient, to decide the approach/equipment required. Access to the NHSGGC Aseptic Non Touch Technique (ANTT®) guideline can be found [here](#).

**Management of care equipment:** All reusable care equipment should be clean at point of use.

Further information can be found [here](#).

**Management of blood and body fluid spillage:** All blood and body fluid spillage should be cleaned.

**Management of waste:** All waste should be disposed of in an appropriate clinical waste bag or sharps bin.

**Occupational risk:** Appropriate procedures and guidance should be followed to reduce the risk of occupational hazards (e.g. needle stick injuries). Needlesafe equipment (e.g. safety hypodermic needles) should be used to reduce the risk of needlestick injury.

**Vascular Access Devices:** All VADs **must** be removed when no longer clinically indicated.

VADs that are not in daily use, e.g. central VADs and dialysis CVCs, may require to be 'locked' to reduce the risk of occlusion. Sterile sodium chloride 0.9% solution should be used to flush and lock VADs. In the event that heparinised saline and taurolidine citrate is used, this must be clearly documented as an exception to normal practice. Consideration should be given to the type and strength of 'lock' that is in place to decide on whether aspiration of 'lock' is required. For example, sodium chloride would not normally require to be aspirated and discarded however, heparin/taurolock **must** be aspirated and discarded to avoid the administration of the 'lock' to the general circulation of the patient as this will significantly increase the risk of coagulopathy.

The practitioner responsible for the clinical area must be informed of:

- any change in the patient's condition
- any change in the VAD insertion site and/or IV in-line pressure monitoring, which cannot be accounted for
- any medication incident or near miss that has occurred. Medical staff must be informed when a medication incident has occurred and a Datix report completed. The patient involved with the medication error/incident must be informed

The VAD and insertion site must be checked for patency and signs of phlebitis before, during and after the administration of any IV flush or medicine.

If the VAD is not being used for continuous infusions, then the patency of the device should be assessed at least once per day (unless the central VAD has been appropriately 'locked' – see [Vascular Access Device guideline](#) for more information).

If the visual inspection phlebitis score (VIP) is 2 or more then the PVC should be removed, documented on PVC care plan/bundle and re sited if required. A PVC/CVC care plan/bundle is required to be completed at least twice per day.

**Needle Free Access Devices (NFAD):** A NFAD is a capless valve that is attached to a VAD. The purpose of NFAD is to reduce the risk of catheter related blood stream infections (CRBSI) and needle stick injuries. These should be attached to VADs, unless these connectors are inappropriate for use with a particular VAD.

All NFADs, whether single or multiple, with or without extension sets, must be primed before use with sterile IV 0.9% sodium chloride. There are a variety of different NFADs available and practitioners should choose a NFAD appropriate to the patient's need and VAD requirements.

#### **Cleaning of access ports/NFAD:**

**When accessing any VAD, care should be taken to adequately clean the access port or NFAD for 30 seconds with a 2% chlorhexidine in 70% isopropyl alcohol wipe ('scrub the hub') and allow to air dry.**

Anti-reflux valve NFADs are also available and should be considered for use when high risk for extravasation medications and therapies are being administered.

Port protectors (alcohol impregnated) may also be considered as part of a strategy to reduce the risk of catheter related blood stream infections (CRBSI). When removing a port protector to connect an administration set or syringe, the NFAD end should still be decontaminated with 2% chlorhexidine in 70% isopropyl alcohol wipe ('scrub the hub') for a minimum of 30secs and allow to dry.

IV medicines/fluids/flush should be administered via the NFAD. The top port of the PVC should **not** be used to routinely administer IV medication in clinical practice. This area cannot be cleaned adequately and therefore the risk of introducing infection is high. The top port of the PVC should **only** be used on sterile (brand new) PVC (immediately following insertion) and in emergency situations.


## 10. General principles

Individual patient condition is fundamental in decision making regarding a suitable method of IV medicine administration e.g. IV bolus administration/intermittent infusion or continuous infusion. Consideration must be given to patient's clinical condition, available VAD and the medicine being administered (the following medicines information resources should be accessed):

- NHSGGC adult and paediatric IV medicine monographs. Electronic copies available via '[clinical info](#)' section via StaffNet or by logging into the Injectable Medicines Guide (username and password are detailed on the StaffNet hub) or contact pharmacy for details.

Clinical areas may also have an IV monograph folder and are responsible for ensuring that these are kept up-to-date by having an allocated person responsible for liaising with pharmacy services

- [British National Formulary](#) (BNF)/[British National Formulary for Children](#) (BNFC)
- [Clinical Drug Information](#) tile on the front page of HEPMA:
- Pharmacy medicine information department or the clinical pharmacist (within working hours – contact via switchboard)
- On call pharmacist (out with working hours - contact via switchboard)
- [NHSGGC Therapeutics Handbook](#)
- The medicine manufacturer's information leaflet
- [Neonatal drug formulary](#) (WoS)



*HEPMA icon*

- One individual patient's IV medicines should be prepared at one time to reduce the risk of error.
- If there is another patient with the same or similar name, a yellow warning sticker should be applied to all prescription records to indicate this (this is different to the yellow labels applied to prepared infusions).
- Practitioners **must** only administer IV medicines that they have either prepared or checked themselves, have been prepared by pharmacy aseptic unit or which are supplied as ready to use preparations (NHSGGC Safe and Secure Handling of Medicines (2023)).
- IV medication should not be prepared and stored in advance of administration, unless there is a clinical rationale, as this increases the risk of bacterial contamination, incorrect medication being administered and may affect stability of the medicine. This should not be normal practice in ward environments. Where interruption to continuous infusion would adversely impact patient safety (e.g. critical care environments), advanced preparation may be necessary. All practitioners involved in the administration of IV medication must be aware of these patient safety risks throughout the process and strategies should be implemented to minimise the occurrence of error.
- Pre prepared solutions/medicines should be used in preference to preparing medicines at ward level e.g. crystalloids with potassium chloride.
- All practitioners should review the need for intravenous therapy on a daily basis and consider IV to oral switchover therapy (IVOST) ([adult IVOST](#); [paediatric IVOST](#)) if appropriate.

- Any change to the IV prescription, resulting in partially filled syringes, bags or bottles must be discarded immediately and not be stored for later use. This fluid/medication is no longer safe to use and must be disposed of following the processes described in the Safe and Secure Handling of Medicines policy (2023).
- Regular monitoring of the infusion site(s) and volume(s) of fluid infused must be recorded using the appropriate documentation, for example VAD care plan/bundle, fluid prescription and fluid balance charts.
- Additional patient monitoring may be required (e.g. ECG monitoring) when administering some IV medications. Consideration should be given to type of IV medication, individual patient clinical condition and duration of infusion.
- Medical staff must be informed when the infusion of medication is not completed due to difficulty with vascular access and this must be documented.
- If a fluid balance chart is required for patient monitoring, it should be available to record the amounts of all fluid volumes infused, with a continuous total maintained for each 24-hour period. This should be documented once the fluids have been administered, rather than when the IV fluid infusion is commenced.
- Frequency of VAD checks by a healthcare practitioner is dependent on clinical area, patient's clinical condition and types of IV medicines being administered e.g. patients receiving irritant medications may have increased frequency checks.
- All IV medicines must be clearly identifiable at all stages during preparation and administration.
- Luer-locking administration sets must be used for all infusion systems. This will include luer-lock syringes for use on syringe devices.
- Anti-reflux valves should be considered when there is a risk of inadvertent backtracking and subsequent medicine overdose when running multiple infusions at different rates.
- When accessing fluid from a rigid container e.g. a glass bottle, the IV administration set will need an air inlet to allow filtered air into the bottle before liquid can easily leave the container. If the air inlet is not integral to the administration set, a separate air inlet should be used. If the liquid is particularly viscous and not easily leaving the container, a sterile air inlet needle which has a filter can be used BUT the administration sets in-line air inlet, if present, must then be put to the closed position as liquid may leak out at this point. This could increase risk of bacterial contamination; liquid could spill onto the electric infusion device and risk both injury and under administration of IV medication.

- During IV medicine administration practitioners should check solutions for signs of crystallization/discolouration/abnormalities.

## 11. Labelling

All infusions prepared with medicines added at ward level must have a yellow IV medicine label completed legibly to identify the following:

- Patient name, CHI number and ward number
- Added medication, LOT number and dose
- Diluent and volume
- Date and time of preparation
- Date and time of expiry for continuous infusions
- Initials of both practitioners involved in the independent 2 person check

The label will be attached securely to the infusion bag/syringe in such a fashion that it does not obscure any other writing/printing or fluid levels. If the label is illegible, smudged, torn or detached from the container, the medicines should not be administered and be discarded. Pharmacy aseptic units will follow their own nationally agreed standards for labelling.

Where multiple infusions are in progress, individual labels identifying the name of medication/infusion fluid must be applied to the infusion line, near the VAD connection point.

When medicines are also being administered via other routes, e.g. IV, epidural and subcutaneous infusions, all lines must be clearly and appropriately labelled with route and medication to avoid inadvertent wrong route errors.

## 12. Priming and changing infusion systems

When an administration set is connected to a VAD, it is essential that a **closed system** is maintained, avoiding unnecessary disconnection. Administration sets are single use and should be discarded once disconnected.

All infusion systems must be fully primed prior to attaching to VAD/NFAD. If an infusion device is being used, the prime function should be utilised.

IV administration sets must be changed every 72 hours. This may be different for certain medications – some administration sets are changed every 12 or 24 hours e.g.

- If the medicine manufacturer's guidance on stability (or the IV monograph) states that the infusion must be changed more frequently than 24 hours



- When parenteral nutrition is being administered
  - When blood/blood products are used. The system should be changed on completion of transfusion or every 12 hours, whichever is soonest
- Check local policy and guidelines.

**All infusion bags and syringes prepared at ward level/clinical area must be changed at least every 24 hours.** Some medications may require the administration set to be changed more frequently. Appropriate documentation should be accurately completed.

Bacterial filters may be used according to local policy and guidelines.

In the unusual circumstance that the closed system is interrupted (e.g. the resiting of a central VAD and administration of an essential medication such as chemotherapy) it is essential to observe a strict aseptic non touch technique to prevent contamination. The practitioner should assess the rationale for the disconnection. The VAD should be flushed with an appropriate volume for the VAD in use of 0.9% sodium chloride (NaCl) in a 10ml syringe, and the administration set sealed with a sterile protective cap. Flush volumes will vary depending on the VAD in use. Flush volume should be large enough to flush the full length of the VAD. Smaller volumes of flush may be considered in neonates and paediatric services.

## 13. Infusion devices and infusion charts

All calculations for device rates should be done **independently** by **both practitioners**. Devices should be programmed and witnessed by **both practitioners**. All device rate changes should be witnessed and verified by **2 practitioners**.

An infusion device should be used to administer IV route medicines and gravity sets only used as an alternative for non-time-critical medications when a device is not available.

An IV infusion prescription record (or pump chart) must be correctly completed by the prescriber and identify the patient and the fluid/medication regime. The volume infused, volume remaining and rate of infusion must be documented hourly on the IV pump chart or equivalent documentation. In paediatric services this information, with addition of a pressure check, will be documented on the Royal Hospital for Children fluid balance chart.

**Infusion devices do not alarm in the event of infiltration or extravasation. Hence, the requirement for site inspection and documentation of VAD insertion site.**

Discontinuation of infusions should be documented on IV infusion chart and the prescription.

### **Syringe device administration**

All syringe devices need to be calibrated to a specific brand of syringe, therefore it is vital to ensure that the correct brand of syringe is used. Incorrect brand of syringe can lead to errors in the rate of the infusion.

Syringe devices should be positioned at a height in line with the patient to reduce the risk of inadvertent free flow.

Luer-lock syringes should be used on syringe devices to reduce the risk of inadvertent disconnection.

When using syringe devices, the flow of fluid can only be seen as being delivered to the patient by confirming the movement of fluid from the syringe to the patient. This will be performed by:

- Cross-referencing the volume delivered and the volume left in the syringe
- Examining the infusion line and any extension set to confirm that they are connected to the patient and that any lumen clamps (of VADs/NFADs) are open to allow the delivery of fluid/medication

### **Volumetric device administration**

When using a volumetric infusion device (not a syringe device), the infusion device shall be confirmed as administering fluid by:

- Observing drops falling within the mini-drop chamber of the infusion line
- Cross-referencing the volume delivered and the volume left in the container (increments are seen at the edge of the printed side of the containers)
- Examining the infusion line and any extension set to confirm that they are connected to the patient and that any lumen clamps (of VADs/NFADs) are open to allow the delivery of fluid/medication

Volume to be infused limits (or drop sensors) must be set on the infusion device whenever possible to minimise the risk of:

- Over or under infusion of fluid/medication
- Air being drawn into the infusion line upon the completion of a reservoir of infusion fluid/medication

## **14. IV flush**

**The fluid administered as an IV flush is sterile sodium chloride 0.9%**

In some instances, additional glucose 5% may be used if it is more suitable for use due to compatibility with the IV medicine being administered. Glucose 5% should also then be flushed

with sterile 0.9% sodium chloride to avoid sticky residue remaining in the VAD and maintain device patency e.g. order of administration being 0.9% sodium chloride (to ensure VAD patency), 5% glucose (to prevent compatibility issues), prescribed medication, 5% glucose (to avoid compatibility issues) then 0.9% sodium chloride (to remove residue and maintain VAD patency).

Situations where IV flush solutions are administered include, but not limited to:

- On insertion of a VAD e.g. PVC or central venous catheter (CVC), to ensure patency
- At specified intervals for patients with a VAD, to maintain device patency
- Before and after the administration of an IV medicine via VAD
- In between each medicine administration when multiple IV medicines are being administered
- Following blood sampling from a VAD, to maintain device patency

Within NHSGGC Adult, Mental Health and Paediatric services (**excluding neonatal and SCBU areas**) it is accepted practice for IV flush solutions to be administered without a formal prescription or administration record as part of the routine care and maintenance of VADs. This is an authorised *exemption* to the normal practice for other Prescription Only Medications (POMs), as described in the NHSGGC Safe and Secure Handling of Medicines (2023) policy. Both sodium chloride 0.9% and glucose 5% injection are classified as POMs due to their intended IV route of administration.

Recommended flush volume is dependent on the VAD or administration set being used and should be a sufficient volume to remove blood or medicine residue. Consider that when the infusion bag is empty the administration set and tubing will still contain an amount of medicine. This will also require to be administered to the patient to ensure the full prescribed dose is administered. Volumetric device and gravity administration sets' generally hold a volume range of 20- 30mL however, this varies depending on the type and length of the set.

Flush volumes are normally twice the internal diameter of the VAD and/or the length (prime volume) of administration set being used.

For PVCs, normal flush volumes would be:

- Adults and children > 1 year            2.5mls - 5mls
- Children ≤ 1 year                        2.5mls

All flushes, regardless of the volume of flush used, should be drawn up using a safety hypodermic needle and syringe which is at least 10ml in size, and administered immediately using a push/pause with positive pressure technique by depressing the syringe plunger before clamping the line or disconnecting the syringe. This syringe size will reduce this risk of excess pressure

being applied on the cannula or vein and/or to prevent VAD fracture/rupture. This technique reduces the risk of backflow of blood into the VAD and/or prevents VAD blockage

All flush solutions should be independently checked with another practitioner prior to administration

**Paediatric patients (under 16 years of age)** require a record of the volume administered to be recorded on the fluid balance chart or other appropriate documentation by the practitioners administering the flush.

Health Care Support Workers (HCSWs) may only administer a flush as part of the PVC insertion process in adults and children > 1 month of age. They must have completed appropriate NHSGGC training and assessed as competent to undertake this skill.

This policy includes flush solutions administered within CT scan environments – in these areas only, local guidelines should be followed with respect to the flush volume/flow rate and size of syringe to use. Within CT areas, no individual flush should exceed 10mls

Student doctors/nurses/midwives may only administer an IV flush as part of the development of clinical skills (e.g. PVC insertion or IV medicine administration) under **direct supervision** by NHSGGC staff who are trained and competent in flush administration during a practice-based placement.

## 15. Review of policy

The policy will be reviewed every 2 years.

## Appendix 1: preparation and administration of an IV bolus

### Equipment requirements:

Prescription chart

PPE (minimum consideration of disposable apron and nitrile gloves)

Clean tray/trolley/appropriate surface

Appropriate size leuc lock syringe(s)

21G safety hypodermic needle/or blunt filter needle

Medication vial(s)/ampoule(s)

IV 0.9% sodium chloride (NaCl) in a 10ml leuc lock syringe(s) – for flush

2% chlorhexidine in 70% isopropyl alcohol wipes

Clinical waste bag

- Perform hand hygiene
- Correctly identify patient, explain process to patient/child +/- parent. Gain consent
- Visually inspect VAD for patency, signs of infiltration, infection or phlebitis
- Perform hand hygiene
- Independently check medicine, reconstitution agent, diluents, flush, prescription and calculation with another practitioner
- Perform hand hygiene
- Apply PPE
- Clean top of vial(s) or ampoule(s) with 2% chlorhexidine and 70% alcohol wipe for 30s. Allow to dry
- Using an aseptic non touch technique, prepare medicine(s) as per the monograph and prescription chart. Label appropriately

### Both practitioners independently undertake

- Scrub the hub of needle free access device (NFAD) for at least 30s using 2% chlorhexidine and 70% alcohol. Allow to dry

*NB \* Aspirate to remove VAD 'lock' if necessary (check patient notes/care plan/bundle). Attach syringe, unclamp lumen if required, aspirate, re apply clamp and detach syringe. Discard.*

- Using an aseptic non touch technique, attach 10ml syringe with 0.9% NaCl to NFAD. Unclamp lumen if required, flush the VAD using a push/pause technique. Reapply clamp and disconnect syringe whilst maintaining positive pressure on syringe plunger

*NB \* Neonates/paediatric may require a smaller volume of flush*

- Flush should be administered with ease. If there are signs of resistance, infiltration, extravasation or inflammation discontinue
- Using an aseptic non touch technique, attach syringe, unclamp lumen if required and administer medicine at a rate in accordance to IV medicine monograph
- If patient complains of discomfort or shows signs of reaction, discontinue and call for help
- Repeat administration process, if multiple bolus administrations are being given, with additional flush between each medicine administration to prevent compatibility issues

**When administration is complete**

- Reposition clamp if required and detach syringe
- Using an aseptic non touch technique, attach 10ml syringe of 0.9% NaCl to NFAD. Unclamp lumen if required, flush VAD using a push/pause technique. Reapply clamp whilst maintaining positive pressure on syringe plunger. Disconnect syringe
- Scrub the hub of needle free access device (NFAD) to remove any medicine residue for at least 30s using 2% chlorhexidine and 70% alcohol. Allow to dry
- Discard all disposable equipment
- Remove PPE and discard as healthcare waste
- Perform hand hygiene
- Both practitioners document medicine administration on appropriate prescription charts

## Appendix 2: preparation and administration of medicines via a volumetric device

### Equipment requirements:

PPE (minimum consideration of disposable apron and nitrile gloves)

Clean tray/trolley/appropriate surface

Appropriate size leuc lock syringe(s)

Medication vial(s)/ampoule(s)

Prescription chart

21G safety hypodermic needle/or blunt filter needle

IV 0.9% sodium chloride (NaCl) in a 10ml leuc lock syringe – for flush

Appropriate volume of diluent (e.g. 100ml 5% glucose)

IV administration set

Yellow medicine additives label

Volumetric device

2% chlorhexidine in 70% isopropyl alcohol wipes

Clinical waste bag

- Perform hand hygiene
- Correctly identify patient, explain process to the patient/child +/- parent. Gain consent
- Visually inspect VAD for patency, signs of infiltration, infection or phlebitis
- Perform hand hygiene
- Independently check medicine, reconstitution agent, diluent, flush, prescription and calculation with another practitioner.
- Complete medicine additive label
- Perform hand hygiene
- Apply PPE
- Clean top of vial(s) or ampoule(s) and injection port of fluid bag (diluent) with 2% chlorhexidine and 70% alcohol wipe for 30s. Allow to dry
- Using an aseptic non touch technique, prepare medicines as per the monograph and prescription chart
- Using an aseptic non touch technique, inject reconstituted medicines slowly into fluid bag (diluent) via additive port. Invert bag 3 to 4 times. Label appropriately

- Using an aseptic non touch technique, attach bag to an appropriate IV administration set and prime line

**Both practitioners independently undertake Chance to Check at bedside**

- Scrub the hub of the needle free access device (NFAD) for at least 30s using 2% chlorhexidine and 70% alcohol. Allow to dry

*NB \* Attach syringe, unclamp lumen if required and aspirate to remove 'lock' if necessary (check patient notes/care plan/bundle). Re apply clamp and detach syringe*

- Using an aseptic non touch technique, attach 10ml syringe with 0.9% NaCl to NFAD. Unclamp lumen if required, flush the VAD using a push/pause technique. Reapply clamp and disconnect syringe whilst maintaining positive pressure on syringe plunger

*NB \* Neonates/paediatric may require a smaller volume of flush*

- Flush should be administered with ease. If there are signs of resistance, infiltration, extravasation or inflammation discontinue process
- Using an aseptic non touch technique, attach the IV administration set to the NFAD, unclamp lumen if required and infuse medicine at an appropriate rate in accordance to IV medicine monographs. A volumetric device should be used where possible
- If patient complains of discomfort or shows signs of a reaction, discontinue and call for help
- Discard all disposable equipment
- Remove PPE and discard as healthcare waste
- Perform hand hygiene
- Both practitioners document medicine administration on appropriate prescription charts (including pump chart) and commence ongoing infusion checks/monitoring

**When administration of medicine is complete**

- Perform hand hygiene
- Apply PPE
- Re position clamp if required

Consider that when the infusion bag is empty the administration set and tubing will still contain an amount of medicine. Volumetric device and gravity administration sets' generally hold a volume range of 20- 30mL, this varies depending on the type and length of the set.

**If disconnecting the administration set:**

- Detach the administration set



- Using an aseptic non touch technique, attach 10ml syringe with 0.9% NaCl to NFAD. Unclamp lumen if required, flush the VAD using a push/pause technique. Reapply clamp and disconnect syringe whilst maintaining positive pressure on syringe plunger.

*NB \* Neonates/paediatric may require a smaller volume of flush*

- Scrub the hub of the needle free access device (NFAD) to remove any medicine residue for at least 30s using 2% chlorhexidine and 70% alcohol. Allow to dry
- Discard all disposable equipment
- Remove PPE and discard as healthcare waste
- Perform hand hygiene
- Both practitioners document medicine administration on appropriate prescription charts (including pump chart) and commence ongoing monitoring

**If leaving the administration set attached to administer the remaining medicine:**

- Using an aseptic non touch technique, remove the completed infusion bag and attach a 50ml or 100ml bag of compatible infusion fluid to the administration set
- Set the device to administer a minimum of 20 – 30mls set at the same rate as the medicine was running at

*NB \* Neonates/paediatric may require a smaller volume of flush*

- Infusion should be administered with ease. If there are signs of resistance, infiltration, extravasation or inflammation discontinue process
- Discard all disposable equipment
- Remove PPE and discard as healthcare waste
- Perform hand hygiene
- Both practitioners document medicine administration on appropriate prescription charts (including pump chart) and commence ongoing monitoring

**When administration is complete**

- Perform hand hygiene
- Apply PPE
- Re position clamp if required
- Detach IV administration set
- Using an aseptic non touch technique, attach 10ml syringe with 0.9% NaCl to NFAD. Unclamp lumen if required, flush the VAD using a push/pause technique. Reapply clamp and disconnect syringe whilst maintaining positive pressure on syringe plunger.

- Discard all disposable equipment
- Remove PPE and discard as healthcare waste
- Perform hand hygiene
- Document administration on appropriate charts (including pump chart) and commence ongoing infusion checks/monitoring

## Appendix 3: preparation and administering medicines via a syringe device

### Requirements

PPE (minimum consideration of disposable apron and nitrile gloves)

Clean tray/trolley/appropriate surface

Appropriate size leur lock syringe(s)

Medication vial(s)/ampoule(s)

Prescription chart

21G safety hypodermic needle/or blunt filter needle

IV 0.9% sodium chloride (NaCl) in a 10ml syringe - for flush

Appropriate volume of diluent (e.g. 0.9% NaCL)

IV infusion line

Yellow medicine additives label

Syringe device

2% chlorhexidine in 70% isopropyl alcohol wipes

Clinical waste bag

- Perform hand hygiene
- Correctly identify patient, explain process to patient/child +/- parent. Gain consent
- Visually inspect VAD for patency, signs of infiltration, infection or phlebitis
- Perform hand hygiene
- Independently check medicine, flush, diluent, prescription and calculation with another practitioner
- Complete medicine additive label
- Perform hand hygiene
- Apply PPE
- Clean top of vial(s) or ampoule(s) for 30s with 2% chlorhexidine and 70% alcohol wipe. Allow to dry
- Using an aseptic non touch technique, prepare medicines as per the monograph and prescription chart
- Using an aseptic non touch technique, draw the reconstituted medicine with any diluents required into the leur lock syringe ensuring that correct end volume is achieved. Label

appropriately

- Attach syringe to an appropriate IV infusion line
- Insert syringe into syringe device, set up according to manufacturer's instruction. Use the syringe device to purge the infusion line of air

**Both practitioners independently undertake Chance to Check at bedside**

- Scrub the hub of the needle free access device (NFAD) for at least 30s using 2% chlorhexidine and 70% alcohol. Allow to dry

*NB \*Attach syringe, unclamp lumen if required and aspirate to remove 'lock' if necessary (check patient notes/care plan/bundle). Re apply clamp and detach syringe*

- Using an aseptic non touch technique, attach 10ml syringe with 0.9% NaCl to NFAD. Unclamp lumen if required, flush the VAD using a push/pause technique. Reapply clamp and disconnect syringe whilst maintaining positive pressure on syringe plunger

*NB \* Neonates/paediatric may require a smaller volume of flush*

- Flush should be administered with ease. If there are signs of resistance, infiltration, extravasation or inflammation discontinue process
- Using an aseptic non touch technique, attach the IV infusion line to the NFAD, unclamp lumen if required and infuse medicine at an appropriate rate in accordance to IV medicine monographs
- If patient complains of discomfort or shows signs of a reaction, discontinue and call for help
- Discard all disposable equipment
- Remove PPE and discard as healthcare waste
- Perform hand hygiene
- Document administration on appropriate charts (including pump chart) and commence ongoing infusion checks and monitoring

**When administration is complete**

- Perform hand hygiene
- Apply PPE
- Re position clamp if required

Consider that when the syringe is empty the infusion line will still contain a small amount of medicine. Syringe infusion lines hold a small volume of solution, generally around 2mL, depending on the type and length of the line. Further consideration should be given as to

whether this small volume should be administered.

**If disconnecting the infusion line:**

- Detach IV infusion line
- Using an aseptic non touch technique, attach 10ml syringe with 0.9% NaCl to NFAD. Unclamp lumen if required, flush the VAD using a push/pause technique. Reapply clamp and disconnect syringe whilst maintaining positive pressure on syringe plunger

*\*Neonates/paediatric may consider a smaller volume of flush*

- Scrub the hub of the needle free access device (NFAD) to remove medicine residue for at least 30s using 2% chlorhexidine and 70% alcohol. Allow to dry
- Discard all disposable equipment
- Remove PPE and discard as healthcare waste
- Perform hand hygiene
- Document administration on appropriate charts (including pump chart) and commence ongoing infusion checks/monitoring

## Appendix 4: Resources

[Safe and secure handing of medicines \(link to be confirmed\)](#)

[Vascular Access Procedure and Practice Guidelines](#)

[Control of Substances Hazardous to Health \(COSHH\)](#)

[Nutrition Resource Manual: Section 5 Part 2 – Parenteral Nutrition](#)

[Infection Prevention and Control](#)

[Sharps – policy and guidance](#)

[Medicine prescription charts](#) - NHSGGC approved in-patient charts

[Injectable medicines guide IV Medusa monographs](#)

[Patient Identification Band Policy](#)

[Medicines Administration Competency Assurance Record](#)

[Consent to Treatment Policy](#)

[Interpreting, Communication, Support and Translation policy](#)

[Aseptic Non Touch Technique \(ANTT®\) \(710\)](#)

[British National Formulary \(BNF\)](#)

[West of Scotland Cancer Network Extravasation in Practice Guidelines \(policy and tools\)](#)

### **Paediatric guidance:**

[Haemato-oncology patient's fluid and electrolytes management \(Schiehallion\)](#)

[Intravenous fluid guidance for previously well children aged 7 days to 16 years](#)

[Intravenous fluid therapy in children and young people in hospital](#)

## Communication and Implementation Plan

Goal: Widespread awareness of updated policy, content, access and supporting resources

Communication Channel	Communication Content	Audience	Timescale/Frequency
Email	<ol style="list-style-type: none"> <li>1. Email ratified policy to END, DNDs, Chief Nurses, Director of Midwifery and Associate Chief Nurses/Midwife for cascade to their teams</li> <li>2. Email ratified policy to Acute Services Lead Nurses and Midwives for cascade to their teams</li> <li>3. Email ratified policy to Acute Services Senior Charge Nurses and Midwives for cascade to their teams</li> </ol>	All healthcare practitioners, nurses, midwives and Operating Department Practitioners who administer IV route medications and flush	TBC once policy ratified
Intranet	<ol style="list-style-type: none"> <li>1. Focus page in Practice Development SharePoint pages</li> </ol>	<ol style="list-style-type: none"> <li>1. Mostly nursing &amp; midwifery staff</li> </ol>	TBC once policy ratified
Virtual Meeting	<ol style="list-style-type: none"> <li>1. Update to Strategic Medicines Improvement Group</li> <li>2. Request agenda item for all Sector/Directorate SUM &amp; ADTC SUM meetings</li> <li>3. Update to Practice Development &amp; Education Forum</li> </ol>	<ol style="list-style-type: none"> <li>1. Medicines safety leaders</li> <li>2. Medicines safety leaders</li> <li>3. All nursing and midwifery educators</li> </ol>	TBC once policy ratified
Education	<ol style="list-style-type: none"> <li>1. Update IV Medicines Administration study day resources</li> <li>2. Add/update/link to any other medicine education resources</li> <li>3. Supporting resources; e.g. ordering details for graphics</li> <li>4. CAS 5 Link Nurse meetings</li> </ol>	<ol style="list-style-type: none"> <li>1. Nurses and midwives</li> <li>2. Nurses and midwives</li> <li>3. All staff</li> <li>4. CAS Link Nurses</li> </ol>	TBC once policy ratified

## Monitoring

Strategies will include, but not be limited to:

- Implementation of the Medicines Safety Audit across Acute Services
- Additional questions on medicine administration are being added to the Combined Care Assurance Audit Tools (CCAAT)
- Review of IV route medicine administration related Datix reports
- Local intelligence and feedback from individuals and forums on medicine related issues.
- Local observation of practice by Senior Charge Nurses/Midwives, Lead Nurses/Midwives and Associate CheifNurses/Midwives

## Impact Assessment

This policy has been reviewed by the Equality and Human Rights Team with no risk of unfair impact on protected characteristics noted.