

NHSGGC Women & Childrens:

Authorised NCIII List for Surgery & Anaesthetics, Paediatrics & Neonates, RHC (Last Updated 18.07.2025):

Department	Designated Lead	Nominated Pharmacy Lead(s)	Designated Contacts within On-Site Pharmacy Department	Nominated Nursing Lead(s)	NCIIIs Approved for Use	Indication	Licensed Status LM / ULM / LM but unlicensed indication	Aseptic Manipulation /Preparation Required	Storage in Clinical Area Authorised	Approved Dose Range	Upper Limit	Folder/ Information Storage Location	Location of Intralipid for Potential LA Toxicity
Paediatric Theatres (Anaesthetics)	Dr Lesley McKee Dr Graham Bell	Susan Kafka Lead Clinical Pharmacist W&C	Nicola Wilson, Senior Pharmacist Paediatric Critical Care Susan Kafka Lead Clinical Pharmacist W&C	Julie Boyle Senior Charge ODP Anaesthetics	Intrathecal levo-bupivacaine ampoules 0.25% and 0.5%	Spinal anaesthesia	Licensed medicine	No. Also used via other routes therefore not subject to NCIII controls within and on leaving pharmacy.	Yes	Up to 1.5mg/kg	Extremely unlikely to exceed 15mg total		Th recovery resus. trolley
					Levo-bupivacaine and clonidine	Spinal anaesthesia	Unlicensed medicine		Yes	Up to 1.5mg/kg for levo-bupivacaine Up to 1microgram/kg for clonidine (always with levo-bupivacaine)	Up to 15mg of levo-bupivacaine Up to 1microgram/kg clonidine (always with levo-bupivacaine)		